

RESEARCH NOTE

Index-based large carnivore population density and abundance estimates for the Ruaha-Rungwa conservation complex in Tanzania

Paolo Strampelli^{1,2}  | Philipp Henschel³ | Charlotte E. Searle^{1,2}  |
Alex Loiruk Lobora⁴  | Halima Kiwango⁵ | David W. Macdonald² | Amy J. Dickman^{1,2}

¹Lion Landscapes, Iringa, Tanzania

²Wildlife Conservation Research Unit, Department of Zoology, Recanati Kaplan Centre, University of Oxford, Oxford, UK

³Panthera, New York, New York, USA

⁴Tanzania Wildlife Research Institute (TAWIRI), Arusha, Tanzania

⁵Tanzania National Parks (TANAPA), Arusha, Tanzania

Correspondence

Paolo Strampelli, Lion Landscapes, Iringa, Tanzania.

Email: paolo.strampelli@gmail.com and paolo.strampelli@zoo.ox.ac.uk

Funding information

Chicago Zoological Society; Columbus Zoo and Aquarium; National Geographic Society; Panthera; Royal Geographical Society; University of Oxford's NERC Environmental Research DTP; The Queen's College of the University of Oxford

Keywords: African wild dog, cheetah, density, leopard, lion, spoor, spotted hyaena, Tanzania

1 | INTRODUCTION

The Ruaha-Rungwa conservation landscape is a ~45,000 km² Protected Area (PA) complex in south-central Tanzania. The area is believed to harbour globally important populations of lion (*Panthera leo*; Bauer et al., 2016), cheetah (*Acinonyx jubatus*; Durant et al., 2017), and African wild dog (*Lycaon pictus*; Woodroffe & Sillero-Zubiri, 2020), as well as important populations of leopard (*Panthera pardus*), spotted hyaena (*Crocuta crocuta*), and striped hyaena (*Hyaena hyaena*) (TAWIRI, 2009). However, no empirical estimates of landscape-wide population abundance exist for these populations, with the only available estimates being based on expert opinion or extrapolations from densities elsewhere (Mesochina et al., 2010; Riggio et al., 2013). Here, we present landscape-level population density and abundance estimates of five large carnivore species in Ruaha-Rungwa. We calculated population density as an index from spoor (track) data, while acknowledging the technique's

low precision (as revealed by recent research) and providing recommendations for future monitoring.

2 | MATERIALS AND METHODS

2.1 | Study area

The complex includes Ruaha National Park (NP), one of the largest NPs in eastern Africa at 20,226 km², which is used for photographic tourism; three Game Reserves (GRs – Rungwa, 9175 km²; Kizigo, 5140 km²; Muhesi, 2720 km²), where trophy hunting tourism is the primary revenue generation mechanism; and a number of multiple-use areas, including Lunda-Mkwambi Game Controlled Area (GCA), Rungwa South Open Area (OA), and MBOMIPA and Waga Wildlife Management Areas (WMAs) (Figure 1). Vegetation cover primarily comprises a mosaic of *Acacia-Commiphora* open savannah/bushland

and central Zambezi and Eastern *Brachystegia*-dominated miombo woodland, complemented by riverine forests and floodplain grasslands (Olson et al., 2001).

2.2 | Study design and data collection

We carried out vehicle-based spoor surveys over two dry seasons, between July and November 2017 and June and November 2018. The study area was divided into 225-km² grid cells, and a minimum of 6 km and a maximum of 20 km were surveyed within all cells with sufficient

road access (Henschel et al., 2020). The maximum 20 km of transects were carried out in all sites where this was possible. For data collection, we employed the protocols presented in Henschel et al. (2020).

2.3 | Spoor and population density estimation

We counted the number of fresh track observations for each species for each transect and from this calculated standardised 'track densities' (Table 1), equivalent to the number of fresh tracks per 100 km of transect (Funston et al., 2010). Population densities were




FIGURE 1 (a) Survey grid and spoor transects in the Ruaha-Rungwa conservation landscape and (b) the landscape within the wider context of Tanzania's protected areas. Both the gazetted and effective boundaries for Ruaha NP are depicted. Rungwa South OA comprises both Rungwa South OA and Rungwa Mzombe OA.

TABLE 1 Survey effort and large carnivore spoor densities in Ruaha-Rungwa

				Spoor density (fresh tracks/100km) ^b				
	Area (km ²)	Spoor surveys (km) ^a	Area coverage	Lion	Leopard	Spotted hyaena	African wild dog	Cheetah
National Parks								
Ruaha NP ^c	17,827	901	72%	8.45 ± 2.62	2.01 ± 1.61	25.81 ± 6.46	3.14 ± 3.35	0.46 ± 0.44
Game Reserves								
Rungwa GR	9175	667	88%	7.60 ± 3.19	8.11 ± 2.15	27.53 ± 5.84	5.32 ± 3.98	0.39 ± 0.76
Kizigo GR	5140	349	85%	8.85 ± 5.74	8.59 ± 3.27	30.92 ± 9.99	0.65 ± 1.28	0.24 ± 0.47
Muhesi GR	2720	278	91%	2.79 ± 2.62	4.22 ± 2.60	19.85 ± 6.74	0.00 ± 0.00	0.00 ± 0.00
Landscape								
All GRs	17,035	1112	86%	6.80 ± 2.47	7.59 ± 1.62	26.81 ± 4.51	3.13 ± 2.28	0.28 ± 0.44
Ruaha NP + GRs	34,862	2013	76%	6.84 ± 1.75	6.76 ± 1.16	25.38 ± 3.83	3.25 ± 2.05	0.38 ± 0.33
Ruaha-Rungwa (inc. GCA & OA)	41,757	2393	85%	6.26 ± 1.58	6.45 ± 1.09	24.43 ± 3.46	2.99 ± 1.82	0.54 ± 0.49

^aSurveys carried out along boundaries of two protected areas are used in calculations for both.

^bCalculated as per Henschel et al. (2020).

^cExcludes permanently settled areas in southwestern Ruaha NP.

TABLE 2 Large carnivore population density and abundance estimates for the different components of the Ruaha-Rungwa complex

	Lion		Leopard		Spotted hyaena		African wild dog		Cheetah		
	Area (km ²)	Density ^a	Population ^b	Density	Population	Density	Population	Density	Population	Density	Population
National Parks											
Ruaha NP ^c	17,827	2.65 (0.00–6.15)	473 (0–1097)	2.01 (0.02–3.99)	358 (3–712)	7.92 (0.00–16.24)	1412 (0–2895)	0.96 (0.00–4.15)	171 (0–741)	0.14 (0.00–0.66)	25 (0–118)
Game Reserves											
Rungwa GR	9175	2.33 (0.00–5.73)	214 (0–526)	2.49 (0.21–4.76)	228 (19–437)	8.45 (2.20–14.69)	775 (202–1348)	1.63 (0.00–6.02)	150 (0–552)	0.12 (0.00–0.89)	11 (0–82)
Kizigo GR	5140	2.71 (0.00–6.71)	139 (0–345)	2.64 (0.45–4.82)	135 (23–248)	9.48 (1.10–17.87)	488 (56–919)	0.20 (0.00–1.14)	10 (0–59)	0.07 (0–0.42)	4 (0–22)
Muhesi GR	2720	0.85 (0.00–2.45)	23 (0–67)	1.29 (0.00–2.88)	35 (0–78)	6.09 (0.71–11.46)	166 (19–312)	N.A.	N.A.	N.A.	N.A.
Rungwa-Kizigo-Muhesi	17,035	2.09 (0.00–5.54)	355 (0–943)	2.33 (0.13–4.52)	397 (23–770)	8.22 (1.47–14.98)	1401 (251–2551)	0.96 (0.00–4.29)	163 (0–730)	0.09 (0.00–0.63)	15 (0–107)
Landscape											
Ruaha-Rungwa (inc. GCA, OA, WMAs)	41,757	1.92 (0.00–5.01)	801 (0–2091)	1.98 (0.00–4.04)	826 (0–1686)	7.50 (0.36–14.63)	3130 (152–6107)	0.92 (0.00–4.21)	382 (0–1756)	0.17 (0.00–1.01)	69 (0–423)

Note: 95% confidence intervals are presented calculated as per Bauer et al. (2017) and Dröge et al. (2020).

^aPopulation density (adult and sub-adult individuals per 100 km²) and associated 95% confidence intervals.

^bEstimated adult and sub-adult population size, using regression equations developed by Funston et al. (2010) and Winterbach et al. (2016) and associated 95% confidence intervals (calculated as per Bauer et al. (2017) and Dröge et al. (2020)).

^cExcludes permanently settled and intensively farmed areas in southwestern Ruaha NP.

then estimated from track densities using the general equation for all large carnivore species developed by Winterbach et al. (2016), modified from Funston et al. (2010). We calculated confidence intervals (CIs) for the population density and abundance estimates using the method employed by Bauer et al. (2017) and Dröge et al. (2020); although this results in wider confidence intervals than the method from Funston et al. (2010), the latter has been shown to overestimate precision (Dröge et al., 2020).

As a result of ongoing land disputes, a portion of southwestern Ruaha NP (~2400 km²) hosts permanent settlements and industrial agriculture activities and was found to be largely devoid of wildlife (Figure 1). Transects in this area were excluded from the analyses, and density and abundance estimates for Ruaha NP refer to the area of the NP without such activities (17,827 km²).

3 | RESULTS AND DISCUSSION

We carried out a total of 2393 km of spoor transects. Overall, we surveyed ~85% of sampling sites in the landscape, ensuring similar sampling in both the primary vegetation types, as well as with regard to distance to rivers, PA boundaries and type, and other natural and anthropogenic features (Figure 1). We detected a total of 151 independent fresh tracks of lion, 149 of leopard, 9 of cheetah, 54 of wild dog, and 572 of spotted hyaena. Table 2 presents overall and PA-specific (where possible) population density and abundance estimates. See Appendix S1 for detection maps for all large carnivores, and Appendix S2 for the track and population density estimates calculations.

Our surveys indicate that Ruaha-Rungwa is home to important populations of lion, leopard, spotted hyaena, and African wild dog. Although cheetah population estimates exhibit very low precision, the low number of detections suggest a relatively sparse population (Table 2). Results confirm that Ruaha-Rungwa is home to an important population of lion, with detections suggesting that the Great Ruaha and Mzombe river valleys are particularly important for the population in the dry season (Appendix S1). Our findings also suggest that Ruaha-Rungwa hosts what is likely to be one of the continent's largest remaining populations of wild dogs, although standard errors for the estimate are especially high due to high variability in spoor densities between transects, which should be kept in mind when interpreting results. Rungwa GR appears particularly important for the species, possibly due to the area consisting primarily of miombo woodlands, which have been suggested to be particularly suitable for the species (Creel, 2001). Spotted hyaena were the species estimated to have the highest abundance in the complex, while leopard track densities were comparable with those from studies employing the same methodology elsewhere (Bauer et al., 2015; Henschel et al., 2020).

Finally, we appreciate that recent research (Dröge et al., 2020) showed that population density and abundance estimates from track densities exhibit lower precision than previously thought.

Indeed, the CIs estimated through the amended formula presented by Dröge et al. (2020) (Table 2) are unlikely to be suitable to monitor population changes over time. Nevertheless, we believe that there is value in sharing our estimates, given the complete lack of empirical landscape-scale population estimates for Ruaha-Rungwa's large carnivores. Going forward, however, we recommend exploring instead the scalability of methods that can provide greater levels of precision, such as spatially explicit capture-recapture (SECR) models applied to data from camera traps (e.g., Strampelli, Henschel, Dickman et al., 2022) or direct sightings (e.g., Elliot & Gopalaswamy, 2016), which have typically been applied over smaller areas. Alternatively, where scalability proves challenging or too resource-intensive, we recommend considering the estimation of alternative robust status parameters (e.g., occupancy; Strampelli, Henschel, Searle et al., 2022) from track data to monitor population status over large scales.

ACKNOWLEDGEMENTS

Fieldwork for this research was carried out under permits 2017-210-NA-2017-107 and 2018-367-NA-2017-107, granted by the Tanzania Commission for Science and Technology (COSTECH) and the Tanzania Wildlife Research Institute (TAWIRI). We would like to thank the Government of Tanzania, TAWIRI, Tanzania National Parks Authority (TANAPA), Tanzania Wildlife Management Authority (TAWA), and Idodi-Pawaga MBOMIPA WMA for their support of this research. We also thank TANAPA Rangers and TAWA and MBOMIPA Game Scouts, as well as Mr. Stivin G. Pangamwene and Mr. Hamis S. Dongo, for their assistance during to fieldwork. We thank Mdonya Old River Camp and Essential Destinations, Ikuka Safari Camp and Nomad Tanzania for their support in Ruaha, and Tanzania Big Game Safaris for their assistance in Rungwa Ikiri. Scholarship funding for PS, and for some fieldwork costs, was provided by the University of Oxford's NERC Environmental Research DTP, and by The Queen's College of the University of Oxford. Additional fieldwork funding was provided by grants from the National Geographic Society, the Columbus Zoological Park Association, Inc., the Chicago Zoological Society, Panthera and the Royal Geographical Society (with IBG), who we thank for their generosity. AD was funded by a Recanati-Kaplan Fellowship.

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare.

DATA AVAILABILITY STATEMENT

Data employed in this study are freely available and can be accessed at: <https://github.com/pstrampelli/RuahaRungwaLargeCarnivoreDensities>.

ORCID

Paolo Strampelli  <https://orcid.org/0000-0003-0763-4164>

Charlotte E. Searle  <https://orcid.org/0000-0002-9861-1429>

Alex Loiruk Lobora  <https://orcid.org/0000-0002-6067-4539>

REFERENCES

- Bauer, H., Henschel, P., Packer, C., Sillero-Zubiri, C., Chardonnet, B., Sogbohossou, E. A., De longh, H. H., & Macdonald, D. W. (2017). Lion trophy hunting in West Africa: A response to Bouché et al. *PLoS One*, 12(3), e0173691.
- Bauer, H., Kamgang, A. S., Kirsten, I., Tumenta, P. N., Saleh, A., Henschel, P., & Sillero-Zubiri, C. (2015). Large carnivore abundance in the Benoue ecosystem, North Cameroon. *African Journal of Ecology*, 54(2), 235–237. <https://doi.org/10.1111/aje.12262>
- Bauer, H., Packer, C., Funston, P., Henschel, P., & Nowell, K. (2016). *Panthera leo* (errata version published in 2017). The IUCN Red List of Threatened Species 2016, 8235. <https://doi.org/10.2305/IUCN.UK.2016-3.RLTS.T15951A107265605.en>
- Creel, S. (2001). Four factors modifying the effect of competition on carnivore population dynamics as illustrated by African wild dogs. *Conservation Biology*, 15(1), 271–274. <https://doi.org/10.1111/j.1523-1739.2001.99534.x>
- Dröge, E., Creel, S., Becker, M. S., Loveridge, A. J., Sousa, L. L., & Macdonald, D. W. (2020). Assessing the performance of index calibration survey methods to monitor populations of wide-ranging low-density carnivores. *Ecology and Evolution*, 10(7), 3276–3292. <https://doi.org/10.1002/ece3.6065>
- Durant, S. M., Mitchell, N., Groom, R., Pettorelli, N., Ipavec, A., Jacobson, A. P., Woodroffe, R., Böhm, M., Hunter, L. T. B., Becker, M. S., Broekhuis, F., Bashir, S., Andresen, L., Aschenborn, O., Beddiaf, M., Belbachir, F., Belbachir-Bazi, A., Berbash, A., de Matos, B., ... Young-Overton, K. (2017). The global decline of cheetah *Acinonyx jubatus* and what it means for conservation. *Proceedings of the National Academy of Sciences*, 114(3), 528–533. <https://doi.org/10.1073/pnas.1611122114>
- Elliot, N. B., & Gopalaswamy, A. M. (2016). Toward accurate and precise estimates of lion density. *Conservation Biology*, 31(4), 934–943. <https://doi.org/10.1111/cobi.12878>
- Funston, P., Frank, L., Stephens, T., Davidson, Z., Loveridge, A., Macdonald, D. M., Durant, S., Packer, C., Mosser, A., & Ferreira, S. M. (2010). Substrate and species constraints on the use of track incidences to estimate African large carnivore abundance. *Journal of Zoology*, 281(1), 56–65. <https://doi.org/10.1111/j.1469-7998.2009.00682.x>
- Henschel, P., Petracca, L. S., Ferreira, S. M., Dennis, S., Laurence, R., & Ekwanga, S. (2020). Census and distribution of large carnivores in the Tsavo national parks, a critical east African wildlife corridor. *African Journal of Ecology*, 58(3), 383–398. <https://doi.org/10.1111/aje.12730>
- Mesochina, P., Mbangwa, O., & Chardonnet, P. (2010). *Conservation status of the lion (Panthera leo Linnaeus, 1758) in Tanzania*. CIRAD.
- Olson, D., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., & D'amico, J. A. (2001). Terrestrial ecoregions of the world: A new map of life on earth. *Bioscience*, 51(11), 933–938. <https://academic.oup.com/bioscience/article-abstract/51/11/933/227116>
- Riggio, J., Jacobson, A., Dollar, L., Bauer, H., Becker, M., Dickman, A., Funston, P., Groom, R., Henschel, P., de longh, H., Lichtenfeld, L., & Pimm, S. (2013). The size of savannah Africa: A lion's (*Panthera leo*) view. *Biodiversity and Conservation*, 22(1), 17–35. <https://doi.org/10.1007/s10531-012-0381-4>
- Strampelli, P., Henschel, P., Dickman, A. J., Searle, C. E., Mkuburo, L., Smit, J. B., Macdonald, D. W., & Ikanda, D. (2022). Camera trapping and spatially explicit capture – Recapture for the monitoring and conservation management of lions: Insights from a globally important population in Tanzania. *Ecological Solutions and Evidence*, 3(1), e12129. <https://doi.org/10.1002/2688-8319.12129>
- Strampelli, P., Henschel, P., Searle, C. E., Macdonald, D. W., & Dickman, A. J. (2022). Habitat use of and threats to African large carnivores in a mixed-use landscape. *Conservation Biology*, e13943. <https://doi.org/10.1111/cobi.13943>
- TAWIRI. (2009). *Tanzania carnivore conservation action plan*. TAWIRI.
- Winterbach, C. W., Ferreira, S. M., Funston, P. J., & Somers, M. J. (2016). Simplified large African carnivore density estimators from track indices. *PeerJ*, 4, e2662. <https://doi.org/10.7717/peerj.2662>
- Woodroffe, R. & Sillero-Zubiri, C. 2020. *Lycaon pictus* (amended version of 2012 assessment). The IUCN red list of threatened species 2020: E.T12436A166502262. <https://doi.org/10.2305/IUCN.UK.2020-1.RLTS.T12436A166502262.en>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Strampelli, P., Henschel, P., Searle, C. E., Lobora, A. L., Kiwango, H., Macdonald, D. W., & Dickman, A. J. (2023). Index-based large carnivore population density and abundance estimates for the Ruaha-Rungwa conservation complex in Tanzania. *African Journal of Ecology*, 61, 221–225. <https://doi.org/10.1111/aje.13092>